



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Equating the values of c as given in (11) and (12),

$$\left(\frac{4x^2(3x-1) - 2x(9x-1)(x+1)}{(9x+1)(x+1)} \right) \frac{d^2y}{dx^2} + \frac{dy}{dx} - \frac{y}{x+1} = 0 \dots (13).$$

$$\therefore 2x(3x^2 + 10x - 1) \frac{d^2y}{dx^2} - (9x+1)(x+1) \frac{dy}{dx} + (9x+1)y = 0,$$

which is the Differential Equation given by the Proposer of the problem.

Scholium.—The proposed Differential Equation is satisfied by the equations, $y=1+x \dots (\alpha)$ and $y=2x^3(3x-1) \dots (\beta)$; that is, these equations are particular solutions, or *particular integrals*, of the Differential Equation.

36. Proposed by H. C. WHITAKER, B. Sc., M. E., Professor of Mathematics, Manual Training School, Philadelphia, Pennsylvania.

A cube is revolved on its diagonal as an axis. Define the figure described and calculate its volume.

II. Solution by the PROPOSER.

The edges adjacent to the axis of revolution generate cones; the two edges not adjacent to the axis generate a hyperboloid of one nappe. Take the axis of revolution as the axis of z . The equation of the cones is $x^2 + y^2 = 2(z \pm \frac{1}{2}\sqrt{3}a)^2$, the altitude of each being $\frac{1}{2}\sqrt{3}a$, the radius of each being $\frac{1}{2}\sqrt{6}a$, the volume of each being $\frac{2\pi\sqrt{3}}{27}a^3$.

The equation of the hyperboloid is $2x^2 + 2y^2 - 4z^2 = a^2$, the volume being the integral of $\pi(2z^2 + \frac{a^2}{2})dz$ between the limits $\frac{1}{2}a\sqrt{3}$ and $-\frac{1}{2}a\sqrt{3}$, this volume being $\frac{5\pi\sqrt{3}}{27}a^3$. Adding the volume of the hyperboloid to the volume of the two cones, the total volume is found to be $\frac{\pi}{\sqrt{3}}a^3 = 1.8138a^3$.

[This solution is given for comparison with that of DR. ZERR, published in last issue. EDITOR.]

38. Proposed by L. B. FILLMAN, St. Petersburg, Pennsylvania.

The diameter of the circular base of a dome is $10=a$ feet, which is also the distance from any point on the circumference of the base to any point on the opposite side of the dome from base to apex. Find the volume of the dome.

I. Solution by GEORGE B. McCLELLAN ZERR, M. A., Ph. D., Professor of Mathematics and Applied Science, Texarkana College, Texarkana, Arkansas-Texas; and the PROPOSER.

Let $x^2 + y^2 = a^2$ be the equation to the circle that forms a section of the dome perpendicular to the base. Then any section parallel to the base at dis-